# CORRECTION Open Access

# Correction: 3-O-acetyl-11-keto-\(\beta\)-boswellic acid exerts anti-tumor effects in glioblastoma by arresting cell cycle at G2/M phase



Wan Li<sup>1,2†</sup>, Jinyi Liu<sup>2,3†</sup>, Weiqi Fu<sup>1,2</sup>, Xiangjin Zheng<sup>1,2</sup>, Liwen Ren<sup>1,2</sup>, Shiwei Liu<sup>4</sup>, Jinhua Wang<sup>1,2\*</sup>, Tengfei Ji<sup>1\*</sup> and Guanhua Du<sup>1,2\*</sup>

# Correction: J Exp Clin Cancer Res 37, 132 (2018) https://doi.org/10.1186/s13046-018-0805-4

Following publication of the original article [1], an error was identified in Fig. 3; specifically:

 Figure 3e and 3f, panels "U251-MG 48h Control" and "U87-MG 48h Control" were inadvertently duplicated

The correction does not have any effect on the results or conclusions of the paper.

## **Author details**

<sup>1</sup>The State Key Laboratory of Bioactive Substance and Function of Natural Medicines, Beijing 100050, China. <sup>2</sup>Key Laboratory of Drug Target Research and Drug Screen, Institute of Materia Medica, Chinese Academy of Medical Science and Peking Union Medical College, Beijing 100050, China. <sup>3</sup>Ethnic Drug Screening & Pharmacology Center, Key Laboratory of Chemistry in Ethnic Medicinal Resources, State Ethnic Affairs Commission & Ministry of Education, Yunnan Minzu University, Kunming 650500, China. <sup>4</sup>Department of Endocrinology, Shanxi DAYI Hospital, Shanxi Medical University, Taiyuan 030002, Shanxi. China.

The original article can be found online at https://doi.org/10.1186/s13046-018-0805-4.

<sup>†</sup>Wan Li and Jinyi Liu contributed equally to this work.

\*Correspondence: wjh@imm.ac.cn; jitf@imm.ac.cn; dugh@imm.ac.cn

Full list of author information is available at the end of the article

Published online: 03 August 2022

### Reference

 Li W, Liu J, Fu W, et al. 3-O-acetyl-11-keto-β-boswellic acid exerts antitumor effects in glioblastoma by arresting cell cycle at G2/M phase. J Exp Clin Cancer Res. 2018;37:132. https://doi.org/10.1186/s13046-018-0805-4.




© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, wist http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

<sup>&</sup>lt;sup>1</sup> The State Key Laboratory of Bioactive Substance and Function of Natural Medicines, Beijing 100050, China

<sup>&</sup>lt;sup>2</sup> Key Laboratory of Drug Target Research and Drug Screen, Institute of Materia Medica, Chinese Academy of Medical Science and Peking Union Medical College, Beijing 100050, China

Li et al. J Exp Clin Cancer Res (2022) 41:236 Page 2 of 2



**Fig. 3** AKBA induces mitochondria-dependent apoptosis in U251 and U87-MG cells in a dose-dependent manner. Flow cytometry using Annexin V-FITC staining shows that AKBA increases apoptosis of U251 (**a**) and U87-MG (**b**) cells. Flow cytometry using JC-1 staining shows that AKBA reduces mitochondrial membrane potential in U251(**c**) and U87-MG (**d**) cells. Flow cytometry using caspase 3/7 live-cell staining shows that AKBA increases the activity of caspase 3/7 in U251 (**e**) and U87-MG (**f**) cells. Western blotting results show that AKBA induces expression of cleaved-caspase 3 and cleaved-PARP in U251 (**g**) and U87-MG (**h**) cells. The experiments were performed in triplicate