CORRECTION Open Access



Correction: Helicobacter pylori CagA promotes epithelial mesenchymal transition in gastric carcinogenesis via triggering oncogenic YAP pathway

Nianshuang Li^{1,2}, Yan Feng², Yi Hu¹, Cong He¹, Chuan Xie¹, Yaobin Ouyang¹, Stephen C. Artim², Degiang Huang¹, Yin Zhu¹, Zhijun Luo³, Zhongming Ge^{2*†} and Nonghua Lu^{1*†}

Correction: *J Exp Clin Cancer Res* 37, 280 (2018) https://doi.org/10.1186/s13046-018-0962-5

Following the publication of the original article [1], duplicate images were found in Figure 7b and Figure 7d. The correct figure is given below:

Published online: 21 November 2023

Reference

 Li N, Feng Y, Hu Y, et al. Helicobacter pylori CagA promotes epithelial mesenchymal transition in gastric carcinogenesis via triggering oncogenic YAP pathway. J Exp Clin Cancer Res. 2018;37:280. https://doi.org/10.1186/ s13046-018-0962-5.

[†]Zhongming Ge and Nonghua Lu contributed equally to this work.

The original article can be found online at https://doi.org/10.1186/s13046-018-0962-5.

*Correspondence: Zhongming Ge zge@mit.edu Nonghua Lu lunonghua@ncu.edu.cn

¹ Department of Gastroenterology, The First Affiliated Hospital of Nanchang University, 17 Yong Waizheng Street, Donghu District, Nanchang 330006, Jiangxi Province, China

 Division of Comparative Medicine, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA
Epartment of Biochemistry, Boston University School of Medicine, 72 East Concord Street, Boston, MA 02118, USA



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativeccommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativeccommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Li et al. J Exp Clin Cancer Res (2023) 42:305 Page 2 of 2




Fig. 7 a, b Wound healing assay (**a**) and Boyden chamber assay (**b**) were performed in AGS cells infected with *H. pylori* wild-type strains (PMSS1 or 7.13) and CagA – mutants. **c** AGS cells were co-cultured with CagA + *H. pylori* strains PMSS1 or 7.13 in combination with VP treatment, subsequently cells invasion was analyzed by transwell assay. **d, e** Cell migration were analyzed by wound healing assay (**d**) and Boyden chamber assay. Data for gene expression are mean ± SEM of 3 independent experiments