

Review

Open Access

Role of retinoic receptors in lung carcinogenesis

Krisztina Bogos*^{1,2,3}, Ferenc Renyi-Vamos^{1,3}, Gabor Kovacs^{2,3}, Jozsef Tovari^{1,3} and Balazs Dome^{1,2,3}

Address: ¹Tumor Biology, Budapest, Hungary, ²Thoracic Oncology National Koranyi Institute of Pulmonology, Budapest, Hungary and ³Department of Tumor Progression, National Institute of Oncology, Budapest, Hungary

Email: Krisztina Bogos* - bogosa@freemail.hu; Ferenc Renyi-Vamos - drrenyi@freemail.hu; Gabor Kovacs - kovac@koranyi.hu; Jozsef Tovari - jtovari@yahoo.com; Balazs Dome - domeb@yahoo.com

* Corresponding author

Published: 14 July 2008

Received: 6 May 2008

Journal of Experimental & Clinical Cancer Research 2008, **27**:18 doi:10.1186/1756-9966-27-18

Accepted: 14 July 2008

This article is available from: <http://www.jeccr.com/content/27/1/18>

© 2008 Bogos et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Several in vitro and in vivo studies have examined the positive and negative effects of retinoids (vitamin A analogs) in premalignant and malignant lesions. Retinoids have been used as chemopreventive and anticancer agents because of their pleiotropic regulator function in cell differentiation, growth, proliferation and apoptosis through interaction with two types of nuclear receptors: retinoic acid receptors and retinoid X receptors. Recent investigations have gradually elucidated the function of retinoids and their signaling pathways and may explain the failure of earlier chemopreventive studies.

In this review we have compiled basic and recent knowledge regarding the role of retinoid receptors in lung carcinogenesis. Sensitive and appropriate biological tools are necessary for screening the risk population and monitoring the efficacy of chemoprevention. Investigation of retinoid receptors is important and may contribute to the establishment of new strategies in chemoprevention for high-risk patients and in the treatment of lung cancer.

Background

Despite antismoking efforts and advances in therapy, lung cancer remains the leading cause of cancer-related death worldwide [1].

Multi-step carcinogenesis has been described as "a gradual accumulation of genetic and epigenetic aberrations resulting in the deregulation of cellular homeostasis" [2]. There is a similarity between bronchial lesions found in carcinogen-treated animals (squamous metaplasia) and the histological changes affecting the bronchial epithelia of humans or animals deficient in vitamin A. Restoration to a normal histological state occurs after vitamin A repl-

tion, which in experimental models has also been shown to confer protection against pro-carcinogens.

The term retinoid (first coined by Sporn in 1976) generally refers to naturally occurring and synthetic vitamin A (retinol) metabolites and analogs [3].

Several studies have shown that vitamin A/retinoids are physiological regulators of embryonic development, vision, reproduction, bone formation, haematopoiesis, differentiation, proliferation and apoptosis. Pharmacologically, they have been recognized as modulators of cell growth, differentiation and apoptosis. Furthermore they

have been shown to suppress carcinogenesis in various organs (e.g. oral cancer, skin, bladder, lung, prostate and breast cancers) in experimental animals [4]. Clinically, retinoids reverse premalignant human epithelial lesions and prevent lung, liver and breast cancer and second primary tumors in the head and neck [5].

It is now generally thought that the effects of retinoids are mainly mediated by the nuclear retinoid receptors, which are members of the steroid and thyroid hormone receptor superfamily [6,7]. Two families of retinoid receptors have been identified, namely RARs and RXRs with three subtypes for each (α ; β , γ) and several isoforms arising from promoter usage and alternate splicing. The retinoid receptors are ligand-activated, DNA binding trans-acting, transcription-modulating proteins. The three RAR types have a strong affinity for all-trans and 9-cis isomers of retinoic acid. The three RXR types, on the other hand, have demonstrated an especially strong specificity for only the 9-cis isomers. Studies have shown that RXR/RAR heterodimers are responsible for transducing the retinoid signal *in vivo* [8]. These heterodimers bind to retinoic acid response elements found in the promoter region of retinoic acid-inducible target genes thereby activating transcription [8,9]. Without ligand RAR-RXR, heterodimers bind to co-repressors, which play an active role in repressing the transcription of targeted genes. The recruitment of histone deacetylases (HDACs) brings about transcriptional repression by preventing the opening of chromatin, which is linked to deacetylation of nucleosomes [10]. Several of the co-activators and co-repressors are shared by multiple signaling pathways, e.g. CBP (cAMP response element binding protein) has been implicated in AP-1 (activator protein 1) and p53 signaling. Meanwhile STAT signaling, Sin3 and HDAC-1 seem to have a role in what Ayer, et. al. call "Mad-Max signaling" [4].

Dawson lists a series of nuclear receptors such as thyroid hormone receptors, vitamin D3 receptors (VDRs), peroxisome proliferator activated receptors (PPARs), and several orphan receptors in which RXR is important as a "heterodimeric partner" [11].

The RARs and RXRs exhibit the conserved module structure of nuclear receptors and their amino acid sequence can be divided into six regions (A-F) based on homology among themselves and with other members of the nuclear superfamily.

The central region C consists of 66 amino acids and has two zinc-binding motifs very much like the core of the DNA binding domain (DBD) which enables cognate response elements to be recognized specifically. Both this central C region and the functionally complex E region are highly conserved between RARs and RXRs. Region E gains

its complexity from the ligand binding domain (LBD), the ligand-dependent transcriptional activation function AF-2, and a dimerization surface contained within it. The name AF-1 has been given to a second transcriptional activation function found in both the amino-terminal A/B regions.

In humans, the genes encoding RAR α , β and γ are respectively located on chromosomes 17q21.1, 3p24 and 12q13. Those for RXR α , β and γ lie on chromosomes 9q34.3, 6p21.3, and 1q22. The physiological importance of the multiple isoforms of RARs is not known precisely, but these isoforms may explain why RARs have pleiotropic biological effects.

There are two major isoforms for RAR α ($\alpha1$ and $\alpha2$) and for RAR γ ($\gamma1$ and $\gamma2$), and four major isoforms for RAR β ($\beta1$ – $\beta4$) and the recently described RAR $\beta1'$ [12], whose absence seems to be responsible for retinoid resistance in lung carcinogenesis. RAR isoforms can be classified as those which are transcribed from either the P1 (class I: RAR $\alpha1$, $\beta1$, and $\beta3$, $\gamma1$) or P2 (class II: RAR $\alpha2$, $\beta2$ and $\beta4$, $\gamma2$) promoter. All class II isoform P2 promoters contain an RA response element and are RA-inducible to varying degrees [13].

Similarly, several isoforms differing from one another in their amino-terminal region have been identified for RXR α ($\alpha1$, $\alpha2$), RXR β ($\beta1$ and $\beta2$), and RXR γ ($\gamma1$, $\gamma2$) [14].

Epigenetic and genetic changes

Respiratory epithelium carcinogenesis is a multifactorial process which includes inherited and acquired genetic changes, chromosomal rearrangements, epigenetic phenomena and chemical carcinogenesis.

Vitamin A deficiency has been associated with bronchial metaplasia and increased lung cancer development. Many other factors contribute to dysfunction of retinoids and their cognate receptors [2].

The first cytogenetic reports connecting chromosome 3 to lung cancers were those of Whang-Peng et al. [15,16], who reported that 100% of small cell lung cancer (SCLC) cases examined showed specific 3p deletions by Giemsa banding. These changes were observed in 12/12 cell lines and three fresh tumors after a two-day culture period. The minimal region of common deletion was 3p14-p25. A number of studies have since been undertaken that obtained similar results that were extended to non small cell lung cancer (NSCLC).

Houle et al. mapped the RAR $\beta2$ at 3p24 and demonstrated that expression was decreased or even suppressed in lung cancer cell lines, suggesting that its re-expression

could suppress malignancy [17]. Frequent loss of RAR β mRNA expression has been described in both primary NSCLCs and bronchial biopsy specimens from heavy smokers [18,19]. Furthermore, in addition to lung cancer [18,20], decreased RAR β 2 mRNA expression has been demonstrated in a variety of solid tumors including head and neck [21] and breast carcinomas [22]. Xu et al. [18] also reported that all RARs and RXRs were expressed in at least 89% of control normal bronchial tissue specimens from patients without a primary lung cancer and that in distant normal bronchus specimens from patients with NSCLC RAR α , RXR α and γ were expressed in more than 95% of the tumor-free specimens. In contrast, RAR β , RAR γ and RXR β expression was decreased, detected in only 76% of NSCLC specimens. Picard et. al similarly showed diminished or absent RAR β protein expression in ~50% of resected NSCLCs [23]. Furthermore, these authors observed normal or elevated RAR α and RXR α expression in NSCLCs. The expression of RAR β , RAR γ , and RXR β was found to be decreased, however, in many tumors, while LOH at 3p24 occurred at a high frequency. This phenomenon was also seen in non-neoplastic lesions. The authors concluded that altered retinoid receptor expression might be involved in lung carcinogenesis. Martinet et al. extended the above study investigating RARs and RXRs alteration in lung cancer precursor lesions. They performed allelotyping for microsatellites located near the RAR/RXR gene loci and immunohistochemistry was additionally carried out to evaluate P53 and RAR β expression. Microsatellite changes occurred frequently in all samples, but without specificity for any group. RAR β marker losses were found in all examined groups, with a concomitant RAR β protein expression [24].

Aberrant methylation of the promoter regions of genes is a major mechanism of gene silencing in tumors [25]. Virmany et al. [26] identified hypermethylation as the underlying mechanism for this frequent loss of RAR β expression. Twenty-one of 49 (43%) primary resected NSCLC samples showed RAR β hypermethylation. In addition, it was demonstrated that RAR β hypermethylation was also important in the pathogenesis of SCLCs, with 62% of SCLCs methylated for RAR β . In the same study, it was also demonstrated that treatment of lung cancer cell lines with the demethylation agent 5-aza-2'-deoxycytidine (5-AZA-CdR) can restore RAR β expression. Moreover, a phase I-II trial in patients with stage IV NSCLC suggests that 5-AZA-CdR may have some clinical activity against metastatic NSCLC [27]. The loss of RAR β mRNA expression has been observed in many lung cancer cell lines also suggesting that to function as a tumor suppressor gene, RAR β expression is contingent on the intracellular concentration of retinoids [28]. The effects of retinol (vitamin A) depend on its intracellular metabolism including its transport by specialized proteins such as Cellular Retinol

Binding Proteins (CRBP) and on its binding as retinoic acid to specific nuclear receptors: the Retinoic Acid Receptors (RARs) and the Retinoid X Receptors (RXRs) [7]. The CRBP I and II transport retinol in the cell and serve as chaperon proteins to prevent unscheduled retinol catabolism. It is the first building block in retinoic acid synthesis.

Retinoid signaling

The mechanisms through which retinoids suppress carcinogenesis, although complex, are gradually being elucidated. Their complexity results from the large number of genes involved in tumor cell differentiation and proliferation that include retinoic acid response elements in their promoters. Retinoids also inhibit tumorigenesis and tumor growth through their ability to induce either apoptosis (programmed cell death) or terminal differentiation. Interestingly, it has been established that the apoptotic process triggered by Retinoid Related Molecules is independent of p53 activation and proceeds through a novel pathway in which the mitochondrion seems to play a pivotal role [29].

As Karamouzis et al. stated in a recent publication [2], a 'switch on/off' model determines the relationship between retinoid receptors and other signaling pathways during bronchial carcinogenesis. According to this model, RXR selective compounds specifically inhibit AP-1 (activator protein 1) activity resulting in inhibited cell proliferation in normal respiratory epithelium, RAR β and RXR α . AP-1-dependent interaction with other nuclear receptors, such as PPAR γ with contribution cofactors (CBP/p300cAMP response element binding protein), ensures cyclin D1 mediated cell cycle inhibition, hence favoring apoptosis or differentiation. Down-regulation of the RAR β mechanism (as detailed above) combined with CBP and AP-1 up-regulation triggers tumor progression and proliferation. Concurrently the inability of RXR α to form heterodimers with PPAR γ enables an AP1/CBP-dependent up-regulation of Cox2, resulting in the inhibition of apoptosis. This crucial role of RXRs may explain the observation of Brabender et. al as well. They observed suppressed mRNA expression of all subtypes of RXRs in curatively resected NSCLC that is followed by statistically worse overall survival [30].

In addition, retinoids play a central role in tumor stroma production and thus in the control of tumor progression and invasion through their ability to regulate the expression of matrix metalloproteinases, transforming growth factor- β , and cell cycle regulator proteins, such as cyclin dependent kinase I, such as p16, or p21 [31,32].

Up to now, the use of retinoids in clinical trials has been limited because of their pharmacologic effects and side effects. Furthermore a majority of human or experimental

NSCLCs are resistant to all *trans*-retinoic acid, and the mechanism of retinoic acid resistance has not been totally elucidated. The absence of the newly recognized RAR β 1' (alternatively spliced from RAR β 1 isoform) could be one reason for retinoid resistance in lung carcinogenesis [12]. In that study RAR β 1' expression was repressed in RA-resistant BEAS-2B-R1 cells in lung cancer, compared with adjacent normal lung tissues. In H358 lung cancer cells that were transiently transfected with RAR β 1', RA treatment was able to restore target gene expression. In order to better understand the mechanism of RAR β 1' repression more studies are needed, and the authors note that "potential reexpression in lung cancer may be important to future approaches to lung cancer chemoprevention" [33].

Chemoprevention

Chemoprevention has been defined as: the application of natural or synthetic molecules to prevent, inhibit or reverse the carcinogenic machinery [34].

For the respiratory tract there are two major classes of agents which appear to prevent damage induced by inhaled carcinogens: retinoids and antioxidants. (In addition to those mentioned above, new classes of chemopreventive agents are under investigation, such as EGFR inhibitors, farnesyl transferase inhibitors, Cyclooxygenase-2 inhibitors etc., but presently we are focused on retinoids and synthetic Retinoid Related Molecules (RRMs).

Clinical trials have shown how complex the chemoprevention approach is. Nevertheless, large primary prevention trials in volunteers (physicians and nurses in the Physicians Health Study) and in high-risk populations (smokers, ex smokers and asbestos workers in the CARET and ATBC studies, and in the more recent EUROSCAN trial) using either beta-carotene, or the combination of beta-carotene and retinyl palmitate and the combination of beta-carotene and alpha-tocopherol have documented

a higher incidence of lung cancer among smoking participants who received beta-carotene [35,36]. Interestingly a negative effect of beta-carotene supplementation has also been observed in experimental animals exposed to cigarette smoke [37]. In the EUROSCAN study, where retinyl palmitate and/or N-acetylcysteine supplementation were used, no beneficial effects on the incidence of second primary cancer and survival were observed. There was one exception for retinal given to workers exposed to asbestosis, which seemed almost protective against mesothelioma development [38]. One possible explanation for the failure and harm seen in the chemoprevention trials could be the procarcinogenic effect of the toxic oxidative carotene metabolites. The oxidative metabolites induce cytochrome P450 enzymes, lowering the serum levels of retinoic acid and down regulating RXR and RAR β . Nicotine by itself inhibits RAR β expression via methylation.

Further randomized, controlled chemoprevention trials designed to test retinoids, β -carotenes or α -tocopherol defined their target population based on smoking history, preneoplastic changes of the bronchial epithelium, or cancer history [39] (Table 1).

In a recent study Lam S. et al. observed that retinol was not effective in the up-regulation of RAR β in lesions with bronchial dysplasia among individuals who continued to smoke [40].

In addition Khurie FR. et al. reported worse prognosis in stage I. lung cancer, which indicated maintenance of RAR β expression and overexpression of RAR β correlated with increased expression of cyclooxygenase-2, an enzyme that contributes to progressive carcinogenesis and is a marker of poor prognosis [41,42].

The gene promoter hypermethylation is a leading cause of gene silencing. The loss of RAR β expression due to hypermethylation is of interest as are bronchial premalignant

Table 1: Lung cancer chemoprevention trials on retinoic acid analogs

Study	Drug	Number of patients	End point	Result	Reference
ATBC	β -Carotene α -Tocopherol	29,133	Lung cancer	Negative/harmful	[57]
CARET	β -Carotene Retinol	18,314	Lung cancer	Negative/harmful	[58]
EUROSCAN	Retinyl-palmitate N-Acetylcystein	2,592	Second primary tumor	Negative	[38]
Physician's Health Study	β -Carotene	22,071	Lung cancer	Negative	[59]
Pastorino et al.	Retinyl-palmitate	307	Second primary tumor	Positive	[35]
Lam et al.	ADT	112	Bronchial dysplasia	Positive	[40]
Kurie et al.	Fenretinide	82	Metaplasia	Negative	[60]
Lippman et al.	13cRA	1,304	Second primary tumor	Negative	[61]

ATBC, α -Tocopherol, β -Carotene Cancer Prevention Study; CARET, Carotene and Retinol Efficacy Test; ADT, anethole dithioethione (5-[p-methoxyphenyl]-1,2-dithiole-3-thione); 13cRA, 13 cis retinoic acid

lesions in developed lung cancer. Recent publications show that hypermethylation of RAR β 2 genes has a different effect on the development of second primary lung cancers (SPLCs) in NSCLCs depending on smoking status. In current smokers, SPLCs developed more frequently when RAR β was unmethylated than when it was hypermethylated. In the case of former smokers it was the opposite. SPLCs were more prevalent in patients with hypermethylated RAR β . Thus, in active smokers, silencing RAR β expression by hypermethylation has a protective effect against the development of SPLCs, whereas in former smokers RAR β expression (unmethylated) appears to be protective. The authors suggested that in current smokers, the continuous high oxygen tension and free radicals induce apoptosis and offer protection from the SPLCs. This apoptosis may be inhibited by retinoic acid if RAR β is expressed on a normal level [43]. These findings explain in part the previous observation that RAR β expression is associated with poor prognosis among patients who are active smokers [44].

Now available are stronger synthetic retinoids that select for an RAR and RXR type without exposing patients to the kind of retinoid toxicity that had previously been observed [45,46]. Aerosolized early on site, these retinoids have been able to reverse the RR deficiency in stabilizing RAR/RXR expression for increased ligand binding to restore normal cellular differentiation [47]. The authors, collaborating with a French research group, conceived and designed an appropriate RR assay in order to measure efficiently the normal bronchial mucosa level of each Retinoid Receptor's mRNA by real time quantitative relative RT-PCR. This method could be useful for screening the RR's status in the damaged bronchial epithelia of the high-risk patient and for monitoring the efficacy of the different Retinoids used as chemopreventive agents [48].

Summary

There is a large body of literature on clinical and preclinical studies using natural retinoids and related compounds for the prevention and the treatment of cancer [49]. The field of lung cancer chemoprevention has been controversial until now. However, there has also been disappointment in extending the therapeutic use of bexarotene (selective RXR agonist) to patients with NSCLC. Although preclinical data and a phase II clinical trial suggested that bexarotene added to platinum based chemotherapy may improve overall survival [50], a subsequent Phase III clinical trial did not bear this out [51,52]. One possible reason is that solid tumors can acquire and develop intrinsic resistance to retinoids during carcinogenesis. The effects of receptor selective retinoids on NSCLC cell lines were examined by Sun et al. According to their findings 8 of the 37 retinoids showed growth-inhibitory activity (IC₅₀ <10 μ M) against at least two of the eight NSCLC cell lines [53].

CD437, a retinoid with some selectivity toward RAR γ , was highly effective [54]. The RXR selective compounds did show growth inhibitory effects when combined with the RAR retinoids. These results indicated that human lung cancer cell lines have a high degree of resistance to synthetic retinoids [55]. Freemantle et al. have summarized the potential mechanisms of Retinoic Acid resistance. Increased P450 catabolism, drug export (P glycoprotein mediated), sequestration of retinoids by CRABs or other proteins, decreased expression of RARs through promoter methylation, persistent histone deacetylation, RAR rearrangement or mutation in the RAR ligand binding domain, and coactivator alteration or alterations downstream of target gene expression may lead to cellular retinoid resistance. This knowledge should aid in predicting those most likely to benefit from retinoid therapy and in developing strategies to optimize single agent or combination retinoid regimens to overcome resistance [56]. The generation of retinoids and rexinoids with restricted selectivity has opened new possibilities for cancer therapy and chemoprevention. It is probable that demethylating and chromatin remodeling agents currently under clinical investigation could be combined with these new retinoids for a better restoration of RR expression.

Authors' contributions

KB and BD conceived of the study and co-wrote the manuscript. FR, GK and JT provided helpful comments and helped write the paper. All authors read and approved the final manuscript.

Acknowledgements

Grant support: Krisztina Bogos was a recipient of the NATO Fellowship (Le Comité Scientifique de l'OTAN, Organisation du Traité de l'Atlantique Nord) in 2003. Balazs Dome is a recipient of the Bolyai Janos Scholarship of the Hungarian Academy of Sciences. Jozsef Tovari is a recipient of the Eotvos Hungarian State Fellowship. Further support: GVOP-2004-3.2.1., KFIIF 2005, OTKA-F68916, OTKA-NK-73082 (B.D.); Hungarian Ministry of Education, NKFP1a-0024-05 (B.D.); OTKA-D048519, OTKA-F046501, OTKA-K76293 (J.To.); "Oveges Jozsef Program" of the Hungarian Agency for Research Fund Management and Research Exploitation (B.D.).

References

1. Smith W, Khuri FR: **The care of the lung cancer patient in the 21st century: a new age.** *Semin Oncol* 2004, **31**:11-15.
2. Karamouzis MV, Papavassiliou AG: **Retinoid receptor cross-talking respiratory epithelium cancer chemoprevention.** *Trends Molecular Medicine* 2005, **1**:10-16.
3. Sporn MB, Dunlop NM, Newton DL, Smith JM: **Prevention of chemical carcinogenesis by vitamin A and its synthetic analogs (retinoids).** *Fed Proc* 1976, **35**(6):1332-1338.
4. Lotan R: **Retinoids and apoptosis: implications for cancer chemoprevention and therapy.** *J Natl Cancer Inst* 1995, **87**:1655-1657.
5. Hansen LA, Sigman CC, Andreola F, Ross SA, Kelloff GJ, De Luca LM: **Retinoids in chemoprevention and differentiation therapy.** *Carcinogenesis* 2000, **7**:1271-1279.
6. Evans RM: **The steroid and thyroid hormone receptor superfamily.** *Science* 1988, **240**:889-895.
7. Chambon P: **A decade of molecular biology of Retinoic Acid.** *Faseb J* 1996, **10**:940-954.

8. Allenby G, Bocquel MT, Saunders M, Kazmer S, Speck J, Rosenberger M, Lovey A, Kastner P, Grippo JF, Chambon P, AA Levin: **Retinoic acid receptors and retinoid X receptors: interactions with endogenous retinoic acids.** *Proc Natl Acad Sci USA* 1993, **90**:30-34.
9. Sun SY, Lotan R: **Retinoids and their receptors in cancer development and chemoprevention.** *Critical Reviews in Oncology/Hematology* 2002, **41**:41-55.
10. Vigushin DM, Coombes RC: **Targeted histone deacetylase inhibition for cancer therapy.** *Curr Cancer Drug Targets* 2004, **2**:205-218.
11. Dawson MI: **Synthetic retinoids and their nuclear receptors.** *Curr Med Chem Anticancer Agents* 2004, **3**:199-230.
12. Petty WJ, Li N, Biddle A: **A novel retinoic acid receptor beta isoform and retinoid resistance in lung carcinogenesis.** *J Natl Cancer Inst* 2005, **97**(22):1645-1651.
13. Evans TR, Kaye SB: **Retinoids: present role and future potential.** *Br J Cancer* 1999, **80**:1-8.
14. Leid M, Kastner P, Chambon P: **Multiplicity generates diversity in the retinoic acid signalling pathways.** *Trends Biochem Sci* 1992, **10**:427-433.
15. Whang-Peng J, Bunn PA Jr, Kao-Shan C, Lee C, Carney DN, Gazdar A, Minna JD: **A nonrandom chromosomal abnormality, del 3p(14-23), in human small cell lung cancer (SCLC).** *Cancer Genet Cytogenet* 1982, **2**:119-134.
16. Whang-Peng J, Kao-Shan CS, Lee EC, Bunn PA, Carney DN, Gazdar AF, Minna JD: **Specific chromosome defect associated with human small-cell lung cancer; deletion 3p(14-23).** *Science* 1982, **215**:181-182.
17. Houle B, Leduc F, Bradley WE: **Implication of RARB in epidermoid (Squamous) lung cancer.** *Genes Chromosomes Cancer* 1991, **5**:358-366.
18. Xu XC, Sozzi G, Lee JS, Lee JJ, Pastorino U, Pilotti S, Kurie JM, Hong WK, Lotan R: **Suppression of retinoic acid receptor beta in non-small-cell lung cancer in vivo: implications for lung cancer development.** *J Natl Cancer Inst* 1997, **89**:624-629.
19. Ayoub J, Jean-François R, Cormier Y, Meyer D, Ying Y, Major P, Desjardins C, Bradley WE: **Placebo-controlled trial of 13-cis-retinoic acid activity on retinoic acid receptor-beta expression in a population at high risk: implications for chemoprevention of lung cancer.** *J Clin Oncol* 1999, **17**(11):3546-3552.
20. Gebert JF, Moghal N, Frangioni JV, Sugarbaker DJ, Neel BG: **High frequency of retinoic acid receptor beta abnormalities in human lung cancer.** *Oncogene* 1991, **6**:1859-6.
21. Xu XC, Ro JY, Lee JS, Shin DM, Hong WK, Lotan R: **Differential expression of nuclear retinoid receptors in normal, premalignant, and malignant head and neck tissues.** *Cancer Res* 1994, **54**:3580-7.
22. Widschwendter M, Berger J, Daxenbichler G, Müller-Holzner E, Widschwendter A, Mayr A, Marth C, Zeimet AG: **Loss of retinoic acid receptor beta expression in breast cancer and morphologically normal adjacent tissue but not in the normal breast tissue distant from the cancer.** *Cancer Res* 1997, **57**:4158-61.
23. Picard E, Seguin C, Monhoven N, Rochette-Egly C, Siat J, Borrelly J, Martinet Y, Martinet N, Vignaud JM: **Expression of retinoid receptor genes and proteins in non-small-cell lung cancer.** *J Natl Cancer Inst* 1999, **91**:1059-66.
24. Martinet N, Alla F, Farré G, Labib T, Drouot H, Vidili R, Picard E, Gaube MP, Le Faou D, Siat J, Borelly J, Vermeylen P, Bazarbachi T, Vignaud JM, Martinet Y: **Retinoic acid receptor and retinoid X receptor alterations in lung cancer precursor lesions.** *Cancer Res* 2000, **60**:2869-2875.
25. Baylin SB, Esteller M, Rountree MR, Bachman KE, Schuebel K, Herman JG: **Aberrant patterns of DNA methylation, chromatin formation and gene expression in cancer.** *Hum Mol Genet* 2001, **10**:687-92.
26. Virmani AK, Rathi A, Zöchbauer-Müller S, Sacchi N, Fukuyama Y, Bryant D, Maitra A, Heda S, Fong KM, Thunnissen F, Minna JD, Gazdar AF: **Promoter methylation and silencing of the retinoic acid receptor-beta gene in lung carcinomas.** *J Natl Cancer Inst* 2000, **92**:1303-1307.
27. Momparler RL, Bouffard DY, Momparler LF, Dionne J, Belanger K, Ayoub J: **Pilot phase I-II study on 5-aza-2'-deoxycytidine (Decitabine) in patients with metastatic lung cancer.** *Anticancer Drugs* 1997, **8**:358-68.
28. Toma S, Emionite L, Fabia G, Spadini N, Vergani L: **Chemoprevention of tumors: the role of RAR-beta.** *Int J Biol Markers* 2003, **18**:78-81.
29. Garattini E, Gianni M, Terao M: **Retinoid related molecules an emerging class of apoptotic agents with promising therapeutic potential in oncology: pharmacological activity and mechanisms of action.** *Curr Pharm Des* 2004, **10**:433-448.
30. Brabender J, Danenberg KD, Metzger R, Schneider PM, Lord RV, Groshen S, Tsao-Wei DD, Park J, Salonga D, Hölscher AH, Danenberg PV: **The role of retinoid X receptor messenger RNA expression in curatively resected non-small cell lung cancer.** *Clin Cancer Res* 2002, **8**:438-443.
31. Ayer DE, Lawrence QA, Eisenman RN: **Mad-Max transcriptional repression is mediated by ternary complex formation with mammalian homologs of yeast repressor Sin3.** *Cell* 1995, **80**:767-776.
32. Hassig CA, Fleischer TC, Billin AN, Schreiber SL, Ayer DE: **Histone deacetylase activity is required for full transcriptional repression by mSin3A.** *Cell* 1997, **89**:341-347.
33. Sabichi AL, Xu X, Lippman SM: **RARBeta1': primed to fight retinoid resistance in lung carcinogenesis.** *J Natl Cancer Inst* 2005, **97**:1632-1633.
34. Krishnan K, Campbell S, Abdel-Rahman F, Whaley S, Stone WL: **Cancer chemoprevention drug targets.** *Curr Drug Targets* 2003, **4**:45-54.
35. Pastorino U, Infante M, Maioli M, Chiesa G, Buysse M, Firket P, Rosmentz N, Clerici M, Soresi E, Valente M: **Adjuvant treatment of stage I lung cancer with high-dose vitamin A.** *J Clin Oncol* 1993, **7**:1216-1222.
36. Pastorini U: **Beta-carotene and the higher risk of lung cancer.** *J Natl Cancer Inst* 1997, **89**:456-457.
37. Wang XD, Liu C, Bronson RT, Smith DE, Krinsky NI, Russell M: **Retinoid signaling and activator protein-1 expression in ferrets given beta-carotene supplements and exposed to tobacco smoke.** *J Natl Cancer Inst* 1999, **91**:60-66.
38. van Zandwijk N, Dalesio O, Pastorino U, de Vries N, van Tinteren : **EUROSCAN, a randomized trial of vitamin A and N-acetylcysteine in patients with head and neck cancer or lung cancer.** *J Natl Cancer Inst* 2000, **92**:977-986.
39. Winterhalder RC, Hirsch FR, Kotantoulas GK, Franklin WA, Bunn PA: **Chemoprevention of lung cancer-from biology to clinical reality.** *Ann Oncol* 2004, **2**:185-196.
40. Lam S, Xu X, Parker-Klein H, Le Riche JC, Macaulay C, Guillaud M, Goldman A, Gazdar AF, Lotan R: **Surrogate end-point biomarker analysis in a retinoid chemoprevention trial in current and former smokers with bronchial dysplasia.** *Int J Oncol* 2003, **23**(6):1607-1613.
41. Khuri FR, Lotan R, Kemp BL, Lippman SM, Wu H, Feng L, Lee JJ, Cooksley CS, Parr B, Chang E, Walsh GL, Lee JS, Hong WK, Xu XC: **Retinoic acid receptor-beta as a prognostic indicator in stage I non-small-cell lung cancer.** *J Clin Oncol* 2000, **15**:2798-2804.
42. Khuri FR, Wu H, Lee JJ, Kemp BL, Lotan R, Lippman SM, Feng L, Hong WK, Xu XC: **Cyclooxygenase-2 overexpression is a marker of poor prognosis in stage I non-small cell lung cancer.** *Clin Cancer Res* 2001, **4**:861-867.
43. Kim JS, Lee H, Kim H, Shim YM, Han J, Park J, Kim DH: **Promoter methylation of retinoic acid receptor beta 2 and the development of second primary lung cancers in non-small-cell lung cancer.** *J Clin Oncol* 2004, **22**:3443-3450.
44. Khuri FR, Lotan R: **Retinoids in lung cancer: friend, foe, or fellow traveler?** *J Clin Oncol* 2004, **22**:3435-3437.
45. Zusi FC, Lorenzi MV, Vivat-Hannah V: **Selective retinoids and retinoids in cancer therapy and chemoprevention.** *Drug Discov Today* 2002, **7**:1165-1174.
46. Dahl AR, Grossi IM, Houchens DP, Scovell LJ, Placke ME, Imondi AR, Stoner GD, De Luca LM, Wang D, Mulshine JL: **Inhaled isotretinoin (13-cis retinoic acid) is an effective lung cancer chemopreventive agent in A/J mice at low doses: a pilot study.** *Clin Cancer Res* 2000, **6**(8):3015-3024.
47. Spinella MJ, Dmitrovsky E: **Aerosolized delivery and lung cancer prevention: pre-clinical models show promise.** *Clin Cancer Res* 2000, **8**:2963-2964.
48. Bogos K, Bonnard L, Noel N, Claudot F, Grosdidier G, Siat J, Vignaud JM, Martinet Y, Martinet M: **Retinoid receptors mRNA levels in**

- normal bronchial mucosa and in lung cancer cell lines. *Lung Cancer* 2004, **45**(Suppl 3): abstract
49. Ortiz MA, Bayon Y, Lopez-Hernandez FJ, Piedrafita FJ: **Retinoids in combination therapies for the treatment of cancer: mechanisms and perspectives.** *Drug Resist Updat* 2002, **5**:162-175.
 50. Edelman MJ, Smith R, Hausner P, Doyle LA, Kalra K, Kendall J, Bedor M, Bisaccia S: **Phase II trial of the novel retinoid, bexarotene, and gemcitabine plus carboplatin in advanced non-small-cell lung cancer.** *J Clin Oncol* 2005, **23**:5774-5778.
 51. Tyagi P: **Bexarotene in combination with chemotherapy fails to prolong survival in patients with advanced non-small-cell lung cancer: results from the SPIRIT I and II trials.** *Clin Lung Cancer* 2005, **7**(1):17-9.
 52. Altucci L, Leibowitz MD, Ogilvie KM, de Lera AR, Gronemeyer H: **RAR and RXR modulation in cancer and metabolic disease.** *Nat Rev Drug Discov* 2007, **10**:793-810.
 53. Sun SY, Yue P, Dawson MI, Shroot B, Michel S, Lamph WW, Heyman RA, Teng M, Chandraratna RA, Shudo K, Hong WK, Lotan R: **Differential effects of synthetic nuclear retinoid receptor-selective retinoids on the growth of human non-small cell lung carcinoma cells.** *Cancer Res* 1997, **57**:4931-4939.
 54. Sun SY, Yue P, Chen X, Hong WK, Lotan R: **The synthetic retinoid CD437 selectively induces apoptosis in human lung cancer cells while sparing normal human lung epithelial cells.** *Cancer Res* 2002, **62**:2430-2436.
 55. Crowe DL: **Receptor selective synthetic retinoids as potential cancer chemotherapy agents.** *Curr Cancer Drug Targets* 2002, **2**:77-86.
 56. Freemantle SJ, Spinella MJ, Dmitrovsky E: **Retinoids in cancer therapy and chemoprevention: promise meets resistance.** *Oncogene* 2003, **22**:7305-7315.
 57. Alpha-tocopherol, Beta-Carotene Cancer Prevention Study Group: **The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers.** *N Engl J Med* 1994, **330**:1029-35.
 58. Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, Keogh JP, Meyskens FL, Valanis B, Williams JH, Barnhart S, Hammar S: **Effects of a combination of beta-carotene and vitamin A on lung cancer and cardiovascular disease.** *N Engl J Med* 1996, **334**:1150-5.
 59. Hennekens CH, Buring JE, Manson JE, Stampfer M, Rosner B, Cook NR, Belanger C, LaMotte F, Gaziano JM, Ridker PM, Willett W, Peto R: **Lack of effect of long term supplementation with beta-carotene on the incidence of malignant neoplasms and cardiovascular disease.** *N Engl J Med* 1996, **334**:1145-9.
 60. Kurie JM, Lee JS, Khuri FR, Mao L, Morice RC, Lee JJ, Walsh GL, Broxson A, Lippman SM, Ro JY, Kemp BL, Liu D, Fritsche HA, Xu X, Lotan R, Hong WK: **N-(4-Hydroxyphenyl)retinamide in the chemoprevention of squamous metaplasia and dysplasia of the bronchial epithelium.** *Clin Cancer Res* 2000, **6**:2973-9.
 61. Lippman SM, Lee JJ, Karp DD, Vokes EE, Benner SE, Goodman GE, Khuri FR, Marks R, Winn RJ, Fry W, Graziano SL, Gandara DR, Okawara G, Woodhouse CL, Williams B, Perez C, Kim HW, Lotan R, Roth JA, Hong WK: **Randomized phase III intergroup trial of isotretinoin to prevent second primary tumors in stage I non-small-cell lung cancer.** *J Natl Cancer Inst (Bethesda)* 2001, **93**:605-18.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

