

RESEARCH

Open Access



ACE2 inhibits breast cancer angiogenesis via suppressing the VEGFa/VEGFR2/ERK pathway

Qi Zhang^{1,3†}, Sihong Lu^{1,2†}, Tianfu Li¹, Liang Yu¹, Yunjian Zhang¹, Huijuan Zeng¹, Xueke Qian⁴, Jiong Bi^{2*} and Ying Lin^{1*}

Abstract

Background: Breast cancer angiogenesis is key for metastasis and predicts a poor prognosis. Angiotensin-converting enzyme 2 (ACE2), as a member of the renin-angiotensin system (RAS), was reported to restrain the progression of hepatocellular carcinoma (HCC) and non-small cell lung cancer (NSCLC) through inhibiting angiogenesis. However, the relationship between ACE2 and breast cancer angiogenesis remains unclear.

Methods: The prognosis and relative gene selection were analysed using the GEPIA, GEO, TCGA and STRING databases. ACE2 expression in breast cancer tissue was estimated by reverse transcription-quantitative polymerase chain reaction (qPCR). Breast cancer cell migration, proliferation and angiogenesis were assessed by Transwell migration, proliferation, tube formation, and wound healing assays. The expression of vascular endothelial growth factor A (VEGFa) was detected by qPCR and Western blotting. The phosphorylation of vascular endothelial growth factor receptor 2 (VEGFR2), mitogen-activated protein kinase 1/2 (MEK1/2), and extracellular signal-regulated protein kinase 1/2 (ERK1/2) was examined by Western blotting. Breast cancer metastasis and angiogenesis in vivo were measured using a zebrafish model.

Results: ACE2 was downregulated in breast cancer patients. Patients with higher ACE2 expression had longer relapse-free survival (RFS). In vitro, ACE2 inhibited breast cancer migration. Meanwhile, ACE2 in breast cancer cells inhibited human umbilical vascular endothelial cell (HUVEC) proliferation, tube formation and migration. In the zebrafish model, ACE2 inhibited breast cancer cell metastasis, as demonstrated by analyses of the number of disseminated foci and the metastatic distance. Neo-angiogenesis was also decreased by ACE2. ACE2 downregulated the expression of VEGFa in breast cancer cells. Furthermore, ACE2 in breast cancer cells inactivated the phosphorylation of VEGFR2, MEK1/2, and ERK1/2 in HUVECs.

Conclusions: Our findings suggest that ACE2, as a potential resistor to breast cancer, might inhibit breast cancer angiogenesis through the VEGFa/VEGFR2/ERK pathway.

Trial registration: Retrospectively registered.

Keywords: Breast cancer, Angiogenesis, ACE2, VEGFa, VEGFR2, ERK

* Correspondence: bjiong@mail.sysu.edu.cn; linyong3@mail.sysu.edu.cn

[†]Qi Zhang and Sihong Lu contributed equally to this work.

²Laboratory of Surgery, The First Affiliated Hospital, Sun Yat-Sen University, No. 58 of Zhongshan 2nd road, Yuexiu district, Guangzhou 510080, China

¹Breast Disease Center, The First Affiliated Hospital, Sun Yat-Sen University, No.58 of Zhongshan 2nd road, Yuexiu district, Guangzhou 510080, China

Full list of author information is available at the end of the article



Background

Tumour angiogenesis refers to the formation of new blood vessels in solid tumours for the supply of nutrients and oxygen, hence promoting the spread of tumour cells [1, 2]. It occurs during tumour development and progression and is a key step in tumour metastasis [3, 4]. The vasculature network around the tumour starts to be actively growing and infiltrative in response to the secretion of pro-angiogenic factors by the tumour, and this process is called the angiogenic switch [5, 6]. The new formed tumour vessels are often twisty, rugged and leaky with discontinuous endothelial cell lining and deficient basement membrane and pericyte coverage. These aberrant morphologies lead to impaired vascular maturation, poor vessel functionality and incoherent tumour perfusion [5, 7, 8]. In addition, various tumour-associated stromal cells and the extracellular matrix constitute the tumour environment to sustain angiogenesis during tumour progression [9, 10]. In breast cancer, the microvessel density (MVD) is a pivotal risk factor for metastasis and a predictor of poor prognosis [11–13]. Thus, breast cancer angiogenesis is a promising diagnostic and therapeutic target that should be investigated [14, 15].

As is well accepted, among the pro-angiogenic factors, vascular endothelial growth factor (VEGF) is the major regulator inducing the sprouting and proliferation of endothelial cells [16] while increasing the permeability of vessels [17, 18]. The secretion of VEGF by tumour cells contributes to neovascularization, which in turn helps the generation and development of cancer [19, 20]. The most functional form of VEGF among its isoforms is VEGF_a, which exerts its angiogenic effects by activating VEGFR2 expressed on endothelial cells [21]. Anti-angiogenic drugs targeting the VEGF pathway are currently used in tumour therapy, particularly in non-small cell lung cancer (NSCLC) [22]. However, these drugs do not exert significant therapeutic effects in breast cancer patients. New targets of anti-angiogenic therapy need to be found.

Angiotensin-converting enzyme 2 (ACE2), a member of the renin-angiotensin system (RAS), plays an important role in the cardiovascular system by converting angiotensin I (AngI) to Ang(1–9) [23] and converting AngII to Ang(1–7) [24, 25]. The products of this enzyme have effects on vasodilation, anti-proliferation and anti-fibrosis [26, 27]. Recently, researchers have found that members of the RAS participate in different biological processes in various tumours. AngII was reported to facilitate tumour migration, proliferation, angiogenesis and metastasis by activating AngII type 1 receptor (AT1R) [28–30], while the activation of AngII type 2 receptor (AT2R) promotes tumour proliferation and angiogenesis in lung cancer [31, 32]. ACE2, as well as Ang(1–7), was reported to inhibit the growth of lung

cancer [33–35] and the metastasis of prostate cancer [36], while it indicates better prognosis in hepatocellular carcinoma [37]. However, the effects of ACE2 on breast cancer angiogenesis remain unknown.

In the present study, we analysed the role of ACE2 in breast cancer prognosis using databases and human tissues, verified the anti-angiogenic effects of ACE2 in vitro and in vivo and then revealed the VEGF_a/VEGFR2/ERK pathway.

Methods

Cell culture

Human breast cancer cell lines (MDA-MB-231, MCF-7, BT-549, ZR-75-30, ZR-75-1, MDA-MB-435, BT-474, MDA-MB-468, and T47D) were purchased from the American Type Culture Collection (ATCC; Manassas, VA, USA) and were maintained in Dulbecco's modified Eagle's medium (DMEM; Gibco, USA) supplemented with 10% foetal bovine serum (FBS; Gibco), 1% penicillin and streptomycin (Gibco).

Human umbilical vascular endothelial cells (HUVECs) were isolated from the umbilical cord and propagated in endothelial cell medium (ECM; ScienCell, USA) supplemented with 10% FBS (ScienCell), 1% endothelial cell growth supplement (ScienCell), and 1% penicillin and streptomycin (ScienCell).

All the cells were cultured in a humidified incubator at 37 °C with 5% CO₂.

Cell transfection

MDA-MB-231 cells overexpressing ACE2 (231-lenti-ACE2) and the matched control cells (231-lenti-Vec) were established using lentivirus carrying the pCMV-ACE2-EGFP-puro plasmid (GeneChem, Shanghai, China). Puromycin (2 µg/ml) was used to select the cells with stable overexpression of ACE2.

MCF-7 cells were transfected with small interfering RNA (siRNA; GenePharma, Shanghai, China) to knock down ACE2 using Lipofectamine RNAiMAX (Invitrogen).

Tube formation assay

The breast cancer cell culture medium was changed to serum-free DMEM medium for 48 h and then was collected, centrifuged and filtered to obtain tumour-conditioned medium (TCM).

The pipet tips and 96-well plates were precooled, and growth-factor-reduced Matrigel (BD, Corning, USA) was thawed overnight prior to the assay. The wells of the 96-well plate were coated with 50 µl of Matrigel and incubated for 30 min. HUVECs were starved in ECM without serum for 24 h prior to the assay, and 2 × 10⁴ HUVECs were seeded on the gel with 200 µl of TCM concentrated 75-fold using an ultrafiltration device (Millipore, USA). The TCM from MCF-7 cells was

supplemented with 1% FBS. The tube formation of HUVECs was observed at different time points during a 12-h experimental period using a microscope.

Cell migration assay

For the tumour cell Transwell migration assay, transfected breast cancer cells were seeded in DMEM without FBS in the upper chamber. The lower chamber was filled with DMEM containing 10% FBS. After 24 h (MDA-MB-231 cells) or 48 h (MCF-7 cells), the upper chamber was washed, fixed, dyed and photographed.

For the HUVEC wound healing assay, HUVECs were cultured to 100% confluence, and vertical scratches were made using a pipet tip. Next, the medium was replaced with TCM. The wounds were photographed every 24 h for 3 days, and the migration distance of the HUVECs was measured.

Cell proliferation assay

Transfected breast cancer cells and HUVECs were seeded in 96-well plates. Once the cells were adherent, the culture medium was replaced with TCM. The CCK8 (Dojindo Laboratories, Japan) assay was then conducted according to the manufacturer's instructions, and the absorbance at a wavelength of 450 nm was read using a microplate reader (Sunrise, Tecan, Austria) every day at the same time for 5 days.

Reverse transcription-quantitative polymerase chain reaction (qPCR)

The total RNA from human breast tissues or cell lines was extracted using the TRIzol reagent (Invitrogen) and subjected to reverse transcription using the PrimeScript RT reagent Kit (Takara, Japan) to synthesize cDNA according to the manufacturer's instructions. The qPCR assay was performed using the LightCycler480 system (Roche, Switzerland) and SYBR Green (Takara). The primer sequences are presented in the Additional file 1: Supporting methods.

Western blotting

The total protein from the cell lines was extracted using RIPA lysis buffer (Beyotime, Shanghai, China) with 1% PMSF (Beyotime), and the concentration was measured using the BCA kit (Beyotime) according to the manufacturer's instructions. The proteins were separated by SDS-PAGE gel electrophoresis and transferred to polyvinylidene fluoride film (PVDF; Millipore). The film was blocked and incubated with different primary antibodies (at the dilutions shown in Additional file 1: Supporting methods) and the IgG-hydrogen peroxide (HRP) secondary antibody. The protein blots were exposed to ECL luminol reagent (Millipore) using the Amersham Imager 600 Image system (AI600, USA).

Zebrafish model

Transfected breast cancer cells labelled with the fluorescent carbocyanine dye Dil (orange colour; GeneCopoeia, USA) were microinjected into the perivitelline cavity of tricaine-anaesthetized Tg (fli1: EGFP) zebrafish embryos at 48 h post-fertilization. The embryos were cultivated in Hank's buffer with 1-phenyl-2-thiourea (PTU) to inhibit pigmentation. The migration of tumour cells in the zebrafish at 0 and 24 h after injection was observed by fluorescence microscopy (Zeiss, German). The vascular structure surrounding the tumour site was photographed using a confocal microscope (LSM780; Zeiss, German).

Human breast cancer sample

All the breast cancer and normal tissues were obtained from patients diagnosed with breast cancer who received breast-conserving or radical surgery without any neoadjuvant therapy at the First Affiliated Hospital of Sun Yat-sen University between 2017 and 2018. The usage of human tissues was approved by the ethics committee of the First Affiliated Hospital of Sun Yat-sen University.

Statistical analysis

The expression analysis was performed using Gene Expression Profiling Interactive Analysis (GEPIA) [38, 39]. The Kaplan-Meier survival analysis was conducted using Kaplan-Meier plotter [40, 41] in the Gene Expression Omnibus (GEO) database and the following datasets: E-MTAB-365 GSE11121, GSE12093, GSE12276, GSE1456, GSE16391, GSE16446, GSE16716, GSE17705, GSE17907, GSE19615, GSE20271, GSE2034, GSE20685, GSE20711, GSE21653, GSE2603, GSE26971, GSE2990, GSE31519, GSE3494, GSE37946, GSE42568, GSE45255, GSE4611, GSE4922, GSE5327, GSE6532, GSE7390, and GSE9195. Gene Ontology (GO) enrichment was analysed to explore the function of ACE2 and related genes. The Cancer Genome Atlas (TCGA) was employed to obtain the heat map of ACE2. The String database and the Cytoscape tool cytoHubba were used to calculate the rank of hub genes. The Kyoto Encyclopedia of Genes and Genomes (KEGG) was used for the pathway analysis. All the bioinformatics analyses were performed using the R2: Genomics Analysis and Visualization Platform [42–44].

All the experiments were repeated three times. The statistical significance was analysed by two-tailed Student's t-test using Prism GraphPad 7.0 software and SPSS 24.0 software, and $*P < 0.05$ was assumed to indicate statistical significance.

Results

ACE2 is downregulated in breast cancer tissue, and this downregulation is associated with worse prognosis


To evaluate the role of ACE2 in breast cancer, we first analysed the expression of ACE2 in breast cancer and

normal tissues using the GEPIA database ($n = 1197$). The level of ACE2 in breast cancer tumour tissue was significantly lower than that in normal tissue (Fig. 1a). We subsequently detected the mRNA expression of ACE2 in breast cancer and paired normal tissues from 29 breast cancer patients. In all the patients, the expression of ACE2 in tumour tissue was obviously lower than that in normal tissue, which was consistent with the GEPIA results (Fig. 1b). The clinical and pathological characteristics of the 29 breast cancer patients were separated into two groups according to the median ACE2 expression value (Additional file 2: Table S1). Additionally, further clinical and pathological data stratified by the ACE2 level were analysed using the TCGA database ($n = 1174$; median cut-off; Additional file 3: Table S2). This indicated that ACE2 might act as an anti-tumour factor in breast cancer. We subsequently constructed the Kaplan-Meier curve of breast cancer using the GEO database ($n = 3951$, best cut-off), which showed that a higher ACE2 level was associated with a longer RFS (Fig. 1c). These results revealed that ACE2 might play a favourable role in breast cancer progression.

ACE2 inhibits breast cancer cell migration

To clarify the role of ACE2 in breast cancer progression, we examined the expression of ACE2 in different breast cancer cell lines (MDA-MB-231, BT-549, MCF-7,

ZR-75-30, ZR-75-1, MDA-MB-435, BT-474, 468, and T47D). A lower level of ACE2 was observed in the breast cancer cell lines with higher metastatic potency, such as MDA-MB-231 cells, whereas a higher expression of ACE2 was detected in the cell lines with weaker metastatic ability, such as MCF-7 cells (Fig. 2a). We subsequently established MDA-MB-231 cell lines stably expressing ACE2 (231-lenti-ACE2) and the control cell lines (231-lenti-Vec). Meanwhile, ACE2 was efficiently knocked down in MCF-7 cells by transfection using three different siRNA sequences (MCF-7-siACE2-1, MCF-7-siACE2-2, and MCF-7-siACE2-3) compared with the expression of ACE2 in the negative control cells (MCF-7-si-NC). The mRNA and protein levels of ACE2 expression in the transfected breast cancer cells were altered as expected (Fig. 2b and c). To estimate the effect of ACE2 on breast cancer cells, we conducted the Transwell migration assay and found that the migration of MDA-MB-231 cells overexpressing ACE2 was decreased compared with that of the control cells. While in MCF-7 cells, the migration ability was enhanced after the knock-down of ACE2 compared with that of the negative control (Fig. 2d). We also performed the CCK8 proliferation assay and found that ACE2 inhibited the proliferation of MCF-7 cells, whereas ACE2 had no effect on the proliferation of MDA-MB-231 cells (Fig. 2e). Thus, ACE2 might inhibit the development of breast cancer in vitro.






Fig. 2 ACE2 inhibits breast cancer cell migration. **(a)** Basal expression of ACE2 in breast cancer cell lines. **(b)** mRNA level of ACE2 in MDA-MB-231 cells overexpressing ACE2 and ACE2-knockdown MCF-7 cells. **(c)** Protein levels of ACE2 in MDA-MB-231 cells overexpressing ACE2 and ACE2-knockdown MCF-7 cells. **(d)** Transwell migration assay of transfected MDA-MB-231 and MCF-7 cells. **(e)** CCK8 proliferation assay of transfected MDA-MB-231 and MCF-7 cells. Microscope magnification: 50 \times . * $P < 0.05$, ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$


ACE2 inhibits breast cancer angiogenesis in vitro

Because ACE2 was reported to have a negative correlation with angiogenesis in HCC [37] and NSCLC [45], we further explored whether ACE2 inhibits breast cancer angiogenesis to prevent breast cancer progression. We performed GO enrichment of ACE2 in breast cancer first and found that ACE2 was obviously related to angiogenesis in breast cancer (Fig. 3a). To validate the results of the GO enrichment analysis, which showed that ACE2 influenced breast cancer angiogenesis, we then collected the TCM from the above-transfected breast cancer cells and performed HUVEC proliferation, tube formation and wound healing assays. The proliferation ability of HUVECs in the TCM from 231-lenti-ACE2 cells was decreased compared with that in the TCM from 231-lenti-Vec cells (Fig. 3b). However, the proliferation of HUVECs in the TCM from ACE2-knockdown MCF-7 cells was increased compared with that in the TCM from MCF-7-negative control cells (Fig. 2b). Additionally, the TCM from 231-lenti-ACE2 cells induced HUVECs to develop fewer and smaller tubes than the TCM from 231-lenti-Vec (Fig. 3c). Additionally, increased tube formation was obtained with HUVECs that were

cultivated in the TCM from ACE2-knockdown MCF-7 cells compared with those that grew in the TCM from MCF-7-si-NC cells (Fig. 3c). Moreover, the migration rate of HUVECs in the TCM from 231-lenti-ACE2 cells was decreased compared with that in the TCM from 231-lenti-Vec cells (Fig. 3d). By contrast, HUVECs migrated for a longer distance in the TCM extracted from ACE2-knockdown MCF-7 cells than in the TCM from negative control cells (Fig. 3d). These results revealed that ACE2 inhibited breast cancer angiogenesis in vitro.

ACE2 inhibits breast cancer metastasis and angiogenesis in a zebrafish model

Next, we established a zebrafish model to explore the effects of ACE2 on breast cancer metastasis and angiogenesis in vivo. The Dil-labelled 231-lenti-Vec and 231-lenti-ACE2 cells (stained in orange colour) were microinjected into the perivitelline space of embryos at 48 h post-fertilization, and the tumour sizes of the two groups of breast cancer cells were even (Fig. 4a). Additionally, equal numbers of the Dil-labelled MCF-7-siACE2-1 and MCF-7-si-NC cells were microinjected (Fig. 4a). The metastasis of the breast




cancer cells in the zebrafish was then observed under a fluorescence microscope. Compared with 231-lenti-Vec cells, the 231-lenti-ACE2 cells resulted in fewer disseminated tumour foci and a shorter metastasis distance (Fig. 4b). Conversely, the ACE2-knockdown MCF-7 cells resulted in a larger number of disseminated tumour foci and a longer metastatic distance than the MCF-7-negative control cells (Fig. 4b).

Furthermore, using a confocal microscope, we observed the degree of angiogenesis around the primary tumour and metastasis sites. Because the vessels of the Tg (fli1: EGFP) zebrafish could glow with a green fluorescence, we found that the number of new formed twisty tumour vessels in the zebrafish injected with 231-lenti-ACE2 cells was less than that found in the zebrafish injected with 231-lenti-Vec cells (Fig. 4c). By contrast, the analysis of the primary tumour and metastasis sites in the zebrafish with MCF-7-siACE2 cells showed the development of more rugged tumour vessels than those found in the zebrafish with MCF-7-si-NC cells (Fig. 4c). Accordingly, the results indicated that ACE2 inhibited breast cancer metastasis and angiogenesis in vivo.

ACE2 inhibits the VEGFa/VEGFR2/ERK pathway to suppress breast cancer angiogenesis

To explore the mechanism through which ACE2 inhibits breast cancer angiogenesis, we analysed the correlation of ACE2 with genes that potentially participate in breast cancer angiogenesis using the TCGA database (Fig. 5a). Based on the intersection of the genes in the heat map, we identified six genes highly enriched in the GO term *angiogenesis* (Fig. 5b). We then calculated the rank of the hub genes using the STRING database and the Cytoscape tool cytoHubba and identified VEGFa as the most plausible mediator of ACE2 and the inhibition of breast cancer angiogenesis (Fig. 5c, Additional file 4: Table S3). Further KEGG pathway analysis revealed 289 pathways that might potentially mediate ACE2 and VEGFa (Fig. 5d, Additional file 5: Table S4). Thus, the findings suggested that VEGFa played a role in the anti-angiogenic effect of ACE2 in breast cancer.

To confirm the results obtained with the databases, we detected the expression of VEGFa in MDA-MB-231 cells overexpressing ACE2 and ACE2-knockdown MCF-7 cells. Consistent with the results of the database analysis, the mRNA and protein levels of VEGFa were declined in



the 231-lenti-ACE2 cells compared with the 231-lenti-Vec cells (Fig. 5e and f). Additionally, the expression of VEGFa was upregulated at both the mRNA and protein levels in the ACE2-knockdown MCF-7 cells compared with the MCF7-siNC cells (Fig. 5e and f). This indicated that VEGFa participated in the ACE2-mediated inhibition of breast cancer angiogenesis, and the underlying mechanism was studied further. It has been reported that extracellular signal-regulated kinase (ERK) signalling is a crucial pathway for the regulation of cell proliferation, differentiation, survival and cell motility in both normal and cancer cells [46–48], and the activation of ERK1/2 stimulates the expression of VEGFa in breast cancer cells [49]. Hence, we detected the phosphorylation level of ERK1/2 in transfected breast cancer cells and found that was decreased in the 231-lenti-ACE2 cells compared with the 231-lenti-Vec cells. However, the phosphorylation level of ERK1/2 in MCF-7-siACE2 cells was increased compared with that in the negative control cells (Fig. 5g).

This revealed that ERK signalling played a role in the process through which ACE2 downregulated VEGFa expression in breast cancer cells.

It has been reported that secreted VEGFa usually combines with VEGFR2 on the endothelial cell membrane to promote angiogenesis [7]. Thus, we cultivated HUVECs in TCM from transfected breast cancer cells and then examined the related protein level of the incubated HUVECs. Accordingly, the phosphorylation level of VEGFR2 in HUVECs cultivated in the TCM from 231-lenti-ACE2 cells was decreased compared with that obtained in the TCM from 231-lenti-Vec cells. However, the TCM extracted from MCF-7-siACE2 cells stimulated the phosphorylation of VEGFR2 in HUVECs compared with that obtained with the TCM from negative control cells. It has been reported that the phosphorylation of VEGFR2 can further activate ERK signalling in endothelial cells. In this signalling process, mitogen-activated protein kinases 1 and 2 (MEK1/2) are phosphorylated




Fig. 5 ACE2 inhibits the VEGFa/VEGFR2/ERK pathway to suppress breast cancer angiogenesis. **(a)** Heat map of the correlation of ACE2 with genes participating in breast cancer angiogenesis. **(b)** UpSet plot of the intersection of angiogenic cytokines and ACE2 in breast cancer. **(c)** KEGG pathway enrichment of ACE2 and potentially related genes. **(d)** Venn diagram showing the overlap between ACE2 and VEGFa genes. **(e)** mRNA level of VEGFa in transfected MDA-MB-231 and MCF-7 cells. **(f)** Protein levels of VEGFa in transfected MDA-MB-231 and MCF-7 cells. **(g)** Phosphorylation level of ERK1/2 in transfected MDA-MB-231 and MCF-7 cells determined by Western blot analysis. **(h)** Western blot analysis of the phosphorylation level of VEGFR2, MEK1/2, and ERK1/2 in HUVECs cultivated for 24 h in the TCM of the transfected tumour cells. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$

and activated, which in turn phosphorylate and activate ERK1/2 [50, 51]. The phosphorylated ERK1/2 is subsequently translocated into the nucleus or activates other kinases and transcription factors to induce cell

proliferation, migration, invasion and differentiation [52, 53]. Hence, we detected the phosphorylation levels of MEK1/2 and ERK1/2 in TCM-cultivated HUVECs and found decreased levels in HUVECs cultivated in the

TCM of 231-lenti-ACE2 cells compared with those in the cells grown in the TCM of 231-lenti-Vec cells. However, the TCM of MCF-7-siACE2 cells induced increased phosphorylation of the two factors compared with the TCM of MCF-7-siNC cells (Fig. 5h). This indicated that the ERK pathway might participate in the anti-angiogenic effect of ACE2. Therefore, ACE2 inhibits breast cancer angiogenesis by the VEGFa/VEGFR2/ERK pathway.


Discussion

Although ACE2 is a member of the RAS family, it also reportedly serves as a potential anti-tumour molecule in various malignant diseases, including lung cancer [33–35], prostate cancer [36] and HCC [37]. In our study, we found that the expression of ACE2 in breast cancer was lower than that in normal tissue, and a lower level of ACE2 in breast cancer patients is related to a worse prognosis, which suggests that ACE2 might play a favourable role in breast cancer. Additionally, in line with a previous study [54], ACE2 inhibited the migration of breast cancer cells while suppressing the proliferation of MCF-7 cells. These evidences suggest that ACE2 acts as a potential tumour suppressor in breast cancer and restrains breast cancer progression.

Angiogenesis, a pivotal process in malignant tumour progression [11, 12], is crucial for tumour metastasis and plays an essential role in breast cancer survival and prognosis [13]. Our data revealed the effect of ACE2 on breast cancer angiogenesis, the crucial step of metastasis. In vitro, ACE2 inhibited the ability of breast cancer cells

to promote HUVEC proliferation, tube formation and migration. And in vivo, ACE2 decreased breast cancer cell-induced neo-angiogenesis in a zebrafish model. Additionally, ACE2 impaired the dissemination and metastasis of breast cancer cells in vivo, which was likely due to the ACE2-induced reduction in the number of tumour vessels and the ACE2-mediated deficiency in the tumour microenvironment. These results imply that ACE2, the metastasis inhibitor, might interrupt breast cancer angiogenesis, which, in turn, disturbs the process of metastasis. This anti-angiogenic function of ACE2 in breast cancer is consistent with that observed in other malignant tumours. For example, in HCC, ACE2 is negatively correlated with CD34 [37], and in NSCLC, ACE2 suppresses angiogenesis and inhibits VEGFa expression [45].

Many pro-angiogenic factors mediate tumour angiogenesis, and among these, VEGFa might play a major role in the process through which ACE2 inhibits breast cancer angiogenesis, as demonstrated through our database analysis. Further qPCR and Western blotting results concordantly showed that ACE2 downregulated VEGFa expression in breast cancer cells, similar to the previous results in NSCLC [45]. Additionally, in our study, ACE2 inhibited the phosphorylation of ERK1/2 in breast cancer cells, which revealed that ERK signalling participated in the ACE2-mediated regulation of the expression of VEGFa. This is consistent with that obtained in a previous study, which showed that the ERK pathway modulated VEGFa [49]. It has been well recognized that



VEGFa from tumour cells combines VEGFR2 on the membrane of neighbouring endothelial cells. This combination phosphorylates and activates VEGFR2, which further phosphorylates and activates the ERK signalling pathway. In the cascade reaction of the ERK pathway, MEK1/2 is phosphorylated and activated, which in turn phosphorylates and activates ERK1/2. This leads to the nuclear translocation or activation of other molecules to promote HUVEC proliferation, differentiation and migration, which finally facilitate angiogenesis [55]. In our study, ACE2 inhibited the phosphorylation of VEGFR2, MEK1/2, and ERK1/2 in HUVECs through the downregulation of VEGFa in breast cancer cells. Therefore, the VEGFa/VEGFR2/ERK pathway might be the mechanism by which ACE2 inhibits breast cancer angiogenesis and metastasis (Fig. 6).

Presently, anti-tumour drugs have already been demonstrated to target the VEGF pathway, and among these drugs, bevacizumab has been widely used clinically for the treatment of multiple cancers, particularly NSCLC [22]. However, this drug cannot significantly improve the survival of breast cancer patients, and the mechanisms underlying this unpleasant result remain controversial. However, ACE2, as an upstream mediator of VEGFa, could inhibit breast cancer angiogenesis and might thus be a promising therapeutic target to postpone the progression of breast cancer. To date, various ACE2 agonists have been designed for different diseases. However, the effects of ACE2 agonists on breast cancer angiogenesis remain unknown, and further study is needed.

Conclusions

We explored the role of ACE2 in breast cancer angiogenesis and found that ACE2 could inhibit breast cancer angiogenesis and metastasis both *in vitro* and *in vivo* and suppress the VEGFa/VEGFR2/ERK pathway. This implied the potential therapeutic value of ACE2 in breast cancer.

Additional files

- Additional file 1:** Supporting methods. (DOCX 19 kb)
- Additional file 2:** Table S1. Clinical and pathological characteristics of breast cancer patients. (DOCX 17 kb)
- Additional file 3:** Table S2. TCGA breast cancer clinical data stratified by the ACE2 expression level. (DOCX 20 kb)
- Additional file 4:** Table S3. Hub gene selection. (DOCX 19 kb)
- Additional file 5:** Table S4. KEGG pathways between ACE2 and VEGFa. (DOCX 32 kb)

Abbreviations

ACE2: Angiotensin-converting enzyme 2; AngI: Angiotensin I; AngII: Angiotensin II; AT1R: AngII type 1 receptor; AT2R: AngII type 2 receptor; ATCC: American Type Culture Collection; BF: Bright field;

DMEM: Dulbecco's modified Eagle's medium; ECM: Endothelial cell medium;; EMT: Epithelial-to-mesenchymal transition; FBS: Foetal bovine serum;; GEO: Gene Expression Omnibus; GEPIA: Gene Expression Profiling Interactive Analysis; GO: Gene Ontology; HCC: Hepatocellular carcinoma; HUVEC: Human umbilical vein endothelial cell; KEGG: Kyoto Encyclopedia of Genes and Genomes; MVD: Microvessel density; NSCLC: Non-small cell lung cancer; PVDF: Polyvinylidene fluoride film; qPCR: Reverse transcription-quantitative polymerase chain reaction; RAS: Renin-angiotensin system; RF: Rhodamine fluorescence; siRNA: small interfering RNA; TCGA: The Cancer Genome Atlas; TCM: Tumour conditioned medium; VEGFa: Vascular endothelial growth factor A; VEGFR2: Vascular endothelial growth factor receptor 2; ERK1/2: Extracellular signal-regulated protein kinase 1/2; MEK1/2: Mitogen-activated protein kinase 1/2; Ang(1–7): Angiotensin (1–7); Ang(1–9): Angiotensin (1–9)

Acknowledgements

We thank Dr. ZW from the Department of Vascular Surgery, the First Affiliated Hospital of Sun Yat-sen University, for providing information regarding angiogenesis-related methods.

Funding

This study was funded by the National Natural Science Foundation of China (81702848, 81602520, 81600336), the Provincial Natural Science Foundation of Guangdong (2017A030313596), the Sun Yat-sen University Clinical Research 5010 Program (2016007) and Young teacher training program of Sun Yat-sen University (17ykpy26).

Availability of data and materials

All the data generated or analysed in this study are included in this published article and its [Additional files](#).

Authors' contributions

QZ and SL conducted the laboratory experiments; TL performed the related bioinformatics analysis; LY and YZ directed and corrected the study; HZ and XQ provided the needed reagents; and JB and YL proposed and designed the research. All the authors approved the manuscript for publication.

Ethics approval and consent to participate

Approval and consent obtained for the use of human tissue were obtained from the Institutional Research Ethics Committee of the First Affiliated Hospital of Sun Yat-sen University.

Approval for all the zebrafish experiments was obtained from Sun Yat-sen University Animal Care and Use Committee of the Zebrafish Model Animal Facility, Institute of Clinical and Translational Research, Sun Yat-sen University.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Breast Disease Center, The First Affiliated Hospital, Sun Yat-Sen University, No.58 of Zhongshan 2nd road, Yuexiu district, Guangzhou 510080, China. ²Laboratory of Surgery, The First Affiliated Hospital, Sun Yat-Sen University, No. 58 of Zhongshan 2nd road, Yuexiu district, Guangzhou 510080, China. ³Guangdong Key Engineering Laboratory for Diagnosis and Treatment of Vascular Disease, The First Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510080, China. ⁴The First Affiliated Hospital, Zhengzhou University, Zhengzhou 450000, China.

Received: 26 January 2019 Accepted: 27 March 2019

Published online: 25 April 2019

References

- Sato Y. Molecular diagnosis of tumor angiogenesis and anti-angiogenic cancer therapy. *Int J Clin Oncol*. 2003;8:200–6.

2. Weis SM, Cheresh DA. Tumor angiogenesis: molecular pathways and therapeutic targets. *Nat Med*. 2011;17:1359–70.
3. Carmeliet P, Jain RK. Principles and mechanisms of vessel normalization for cancer and other angiogenic diseases. *Nat Rev Drug Discov*. 2011;10:417–27.
4. Carmeliet P, Jain RK. Angiogenesis in cancer and other diseases. *Nature*. 2000;407:249–57.
5. Bergers G, Benjamin LE. Tumorigenesis and the angiogenic switch. *Nat Rev Cancer*. 2003;3:401–10.
6. Inoue M, Hager JH, Ferrara N, Gerber HP, Hanahan D. VEGF-A has a critical, nonredundant role in angiogenic switching and pancreatic beta cell carcinogenesis. *Cancer Cell*. 2002;1:193–202.
7. Potente M, Gerhardt H, Carmeliet P. Basic and therapeutic aspects of angiogenesis. *Cell*. 2011;146:873–87.
8. Morikawa S, Baluk P, Kaidoh T, Haskell A, Jain RK, McDonald DM. Abnormalities in pericytes on blood vessels and endothelial sprouts in tumors. *Am J Pathol*. 2002;160:985–1000.
9. Bergers G, Brekken R, McMahon G, Vu TH, Itoh T, Tamaki K, et al. Matrix metalloproteinase-9 triggers the angiogenic switch during carcinogenesis. *Nat Cell Biol*. 2000;2:737–44.
10. Egeblad M, Nakasone ES, Werb Z. Tumors as organs: complex tissues that interface with the entire organism. *Dev Cell*. 2010;18:884–901.
11. Carmeliet P, Jain RK. Molecular mechanisms and clinical applications of angiogenesis. *Nature*. 2011;473:298–307.
12. Bielenberg DR, Zetter BR. The contribution of angiogenesis to the process of metastasis. *Cancer J*. 2015;21:267–73.
13. Weidner N, Semple JP, Welch WR, Folkman J. Tumor angiogenesis and metastasis—correlation in invasive breast carcinoma. *N Engl J Med*. 1991;324:1–8.
14. van Hinsbergh VW, Collen A, Koolwijk P. Angiogenesis and anti-angiogenesis: perspectives for the treatment of solid tumors. *Ann Oncol*. 1999;10(Suppl 4):60–3.
15. Pietras RJ, Weinberg OK. Antiangiogenic steroids in human cancer therapy. *Evid Based Complement Alternat Med*. 2005;2:49–57.
16. Tammela T, Enholm B, Alitalo K, Paavonen K. The biology of vascular endothelial growth factors. *Cardiovasc Res*. 2005;65:550–63.
17. Kevil CG, Payne DK, Mire E, Alexander JS. Vascular permeability factor/vascular endothelial cell growth factor-mediated permeability occurs through disorganization of endothelial junctional proteins. *J Biol Chem*. 1998;273:15099–103.
18. Behzadian MA, Windsor LJ, Ghaly N, Liou G, Tsai NT, Caldwell RB. VEGF-induced paracellular permeability in cultured endothelial cells involves urokinase and its receptor. *FASEB J*. 2003;17:752–4.
19. Yoshiji H, Harris SR, Thorgerirsson UP. Vascular endothelial growth factor is essential for initial but not continued *in vivo* growth of human breast carcinoma cells. *Cancer Res*. 1997;57:3924–8.
20. Giavazzi R, Sennino B, Coltrini D, Garofalo A, Dossi R, Ronca R, et al. Distinct role of fibroblast growth factor-2 and vascular endothelial growth factor on tumor growth and angiogenesis. *Am J Pathol*. 2003;162:1913–26.
21. Ferrara N. Vascular endothelial growth factor: molecular and biological aspects. *Curr Top Microbiol Immunol*. 1999;237:1–30.
22. Sandler A, Gray R, Perry MC, Brahmer J, Schiller JH, Dowlati A, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med*. 2006;355:2542–50.
23. Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, et al. A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ Res*. 2000;87:E1–9.
24. Keidar S, Kaplan M, Gamlie-Lazarovich A. ACE2 of the heart: from angiotensin I to angiotensin (1-7). *Cardiovasc Res*. 2007;73:463–9.
25. Wang W, McKinnie SM, Farhan M, Paul M, McDonald T, McLean B, et al. Angiotensin-converting enzyme 2 metabolizes and partially inactivates pyrilapelin-13 and apelin-17: physiological effects in the cardiovascular system. *Hypertension*. 2016;68:365–77.
26. Deshayes C, Nahmias C. Angiotensin receptors: a new role in cancer? *Trends Endocrinol Metab*. 2005;16:293–9.
27. Santos RA, Ferreira AJ, Simoes ESAC. Recent advances in the angiotensin-converting enzyme 2-angiotensin (1-7)-mas axis. *Exp Physiol*. 2008;93:519–27.
28. Egami K, Murohara T, Shimada T, Sasaki K, Shintani S, Sugaya T, et al. Role of host angiotensin II type 1 receptor in tumor angiogenesis and growth. *J Clin Invest*. 2003;112:67–75.
29. Fujita M, Hayashi I, Yamashina S, Itoman M, Majima M. Blockade of angiotensin AT1a receptor signaling reduces tumor growth, angiogenesis, and metastasis. *Biochem Biophys Res Commun*. 2002;294:441–7.
30. Fujita M, Hayashi I, Yamashina S, Fukamizu A, Itoman M, Majima M. Angiotensin type 1a receptor signaling-dependent induction of vascular endothelial growth factor in stroma is relevant to tumor-associated angiogenesis and tumor growth. *Carcinogenesis*. 2005;26:271–9.
31. Kanehira T, Tani T, Takagi T, Nakano Y, Howard EF, Tamura M. Angiotensin II type 2 receptor gene deficiency attenuates susceptibility to tobacco-specific nitrosamine-induced lung tumorigenesis: involvement of transforming growth factor-beta-dependent cell growth attenuation. *Cancer Res*. 2005;65:7660–5.
32. Pickel L, Matsuzuka T, Doi C, Ayuzawa R, Maurya DK, Xie SX, et al. Overexpression of angiotensin II type 2 receptor gene induces cell death in lung adenocarcinoma cells. *Cancer Biol Ther*. 2010;9:277–85.
33. Ni L, Feng Y, Wan H, Ma Q, Fan L, Qian Y, et al. Angiotensin-(1-7) inhibits the migration and invasion of A549 human lung adenocarcinoma cells through inactivation of the PI3K/Akt and MAPK signaling pathways. *Oncol Rep*. 2012;27:783–90.
34. Menon J, Soto-Pantoja DR, Callahan MF, Cline JM, Ferrario CM, Tallant EA, et al. Angiotensin-(1-7) inhibits growth of human lung adenocarcinoma xenografts in nude mice through a reduction in cyclooxygenase-2. *Cancer Res*. 2007;67:2809–15.
35. Qian YR, Guo Y, Wan HY, Fan L, Feng Y, Ni L, et al. Angiotensin-converting enzyme 2 attenuates the metastasis of non-small cell lung cancer through inhibition of epithelial-mesenchymal transition. *Oncol Rep*. 2013;29:2408–14.
36. Krishnan B, Torti FM, Gallagher PE, Tallant EA. Angiotensin-(1-7) reduces proliferation and angiogenesis of human prostate cancer xenografts with a decrease in angiogenic factors and an increase in sFlt-1. *Prostate*. 2013;73:60–70.
37. Ye G, Qin Y, Lu X, Xu X, Wu C, et al. The association of renin-angiotensin system genes with the progression of hepatocellular carcinoma. *Biochem Biophys Res Commun*. 2015;459:18–23.
38. GEPIA. Interactive bodymap. Peking University: The median expression of tumor and normal samples in bodymap. <http://gepia.cancer-pku.cn/detail.php>. Accessed 27 Dec 2018
39. Tang Z, Li C, Kang B, Gao G, Li C, Zhang Z. GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res*. 2017;45:W98–102.
40. Lanczky A, Nagy A, Bottai G, Munkacsy G, Szabo A, Santarpia L, et al. miRpower: a web-tool to validate survival-associated miRNAs utilizing expression data from 2178 breast cancer patients. *Breast Cancer Res Treat*. 2016;160:439–46.
41. Kaplan Meier Plotter. What is the KM plotter? In: Hungarian academy of sciences; 2018. <http://kmplot.com>. Accessed 27 Dec 2018.
42. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*. 2003;13:2498–504.
43. Chin CH, Chen SH, Wu HH, Ho CW, Ko MT, Lin CY. cytoHubba: identifying hub objects and sub-networks from complex interactome. *BMC Syst Biol*. 2014;8(Suppl 4):S11.
44. R2: genomics analysis and visualization platform. Academic Medical Center (AMC) Amsterdam, The Netherlands. 2018. <http://r2.amc.nl>. Accessed 23 Dec 2018.
45. Feng Y, Ni L, Wan H, Fan L, Fei X, Ma Q, et al. Overexpression of ACE2 produces antitumor effects via inhibition of angiogenesis and tumor cell invasion *in vivo* and *in vitro*. *Oncol Rep*. 2011;26:1157–64.
46. Bromberg-White JL, Andersen NJ, Duesbery NS. MEK genomics in development and disease. *Brief Funct Genomics*. 2012;11:300–10.
47. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. 2000;100:57–70.
48. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144:646–74.
49. Liu H, Tang L, Li X, Li H. Triptolide inhibits vascular endothelial growth factor-mediated angiogenesis in human breast cancer cells. *Exp Ther Med*. 2018;16:830–6.
50. Caunt CJ, Sale MJ, Smith PD, Cook SJ. MEK1 and MEK2 inhibitors and cancer therapy: the long and winding road. *Nat Rev Cancer*. 2015;15:577–92.
51. Samatar AA, Poulikakos PI. Targeting RAS-ERK signalling in cancer: promises and challenges. *Nat Rev Drug Discov*. 2014;13:928–42.
52. Pang X, Yi Z, Zhang X, Sung B, Qu W, Lian X, et al. Acetyl-11-keto-beta-boswellic acid inhibits prostate tumor growth by suppressing vascular endothelial growth factor receptor 2-mediated angiogenesis. *Cancer Res*. 2009;69:5893–900.
53. Leelahavanichkul K, Amornphimoltham P, Molinolo AA, Basile JR, Koontongkaew S, Gutkind JS. A role for p38 MAPK in head and neck cancer

cell growth and tumor-induced angiogenesis and lymphangiogenesis. *Mol Oncol.* 2014;8:105–18.

54. Yu C, Tang W, Wang Y, Shen Q, Wang B, Cai C, et al. Downregulation of ACE2/Ang-(1-7)/mas axis promotes breast cancer metastasis by enhancing store-operated calcium entry. *Cancer Lett.* 2016;376:268–77.
55. Kerbel RS. Tumor angiogenesis. *N Engl J Med.* 2008;358:2039–49.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

