

CORRECTION

Open Access

Correction to: The interplay between HIF-1 α and noncoding RNAs in cancer



Xiaofeng Peng^{1,2†}, Han Gao^{3†}, Rui Xu⁴, Huiyu Wang¹, Jie Mei^{1*} and Chaoying Liu^{1*}

Correction to: J Exp Clin Cancer Res

<https://doi.org/10.1186/s13046-020-1535-y>

In the original publication of this manuscript [1], Fig. 2 contains incorrect labels and feedback loops. The revised version of Fig. 2 is shown below.

In the first paragraph of the 'Negative feedback loop between HIF-1 α and ncRNA' section, there are two instances of 'miR-439'; these should instead read 'miR-429'.

Author details

¹Department of Oncology, Wuxi People's Hospital Affiliated to Nanjing Medical University, 299 Qingyang Road, Wuxi 214023, China. ²The First Clinical Medicine School, Nanjing Medical University, Nanjing 211166, China. ³Wuxi School of Medicine, Jiangnan University, Wuxi 214122, China. ⁴School of Basic Medical Sciences, Nanjing Medical University, Nanjing 211166, China.

Published online: 03 March 2020

Reference

1. Peng, et al. *J Exp Clin Cancer Res.* 2020;39:27.

The original article can be found online at <https://doi.org/10.1186/s13046-020-1535-y>

* Correspondence: meijie1996@njmu.edu.cn; liuchaoying666@163.com

[†]Xiaofeng Peng and Han Gao contributed equally to this work.

¹Department of Oncology, Wuxi People's Hospital Affiliated to Nanjing Medical University, 299 Qingyang Road, Wuxi 214023, China

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.




Fig. 2 Reciprocal feedback loops between HIF-1 α and ncRNAs. In addition to a unidirectional regulation pattern, there are several direct or indirect feedback loops between HIF-1 α and ncRNAs. It seems quite feasible that the ncRNAs, HIF-1 α and other co-operators would eventually intertwine to form mutually reciprocal feedback loops in both positive and negative manners. In addition to common feedback loops, lincRNA-p21 and HISLA can block VHL- and PHD-dependent HIF-1 α repression instead of directly interacting with HIF-1 α and other co-operators