

CORRECTION

Open Access



Correction to: Genistein inhibits stemness of SKOV3 cells induced by macrophages co-cultured with ovarian cancer stem-like cells through IL-8/STAT3 axis

Yingxia Ning^{1,2†}, Weifeng Feng^{2†}, Xiaocheng Cao^{4,5}, Kaiqun Ren^{4,5}, Meifang Quan^{4,5}, A. Chen^{4,5}, Chang Xu^{4,5}, Yebei Qiu^{4,5}, Jianguo Cao^{4,5*}, Xiang Li^{3,4,5*} and Xin Luo^{2*}

Correction to: J Exp Clin Cancer Res 38, 19 (2019)
<https://doi.org/10.1186/s13046-018-1010-1>

Following publication of the original article [1], the authors identified minor errors in image-typesetting in Fig. 9; specifically in the panels shown in Fig. 9a and d as follows:

- Figure 9a: a ruler has been added alongside the fragments of the tumor
- Figure 9d: corrected the panel displaying HE staining, GEN (50 μ M) (-) and Ad-shSTAT3 (+) (*top row, middle-right*)

The correction does not have any effect on the results or conclusions of the paper. The original article has been corrected.

Author details

¹Department of Gynaecology and Obstetrics, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou 510120, China. ²The First Affiliated Hospital of Jinan University, Guangzhou 510632, China. ³Department of preclinical medicine, Medical College, Hunan Normal University, Changsha 410013, China. ⁴Department of Pharmaceutical Science, Medical College, Hunan Normal University, Changsha 410013, China. ⁵Key Laboratory of Study and Discover of Small Targeted Molecules of Hunan Province, Changsha 410013, China.

Published online: 04 May 2021

Reference

1. Ning Y, Feng W, Cao X, Ren K, Quan M, Chen A, et al. Genistein inhibits stemness of SKOV3 cells induced by macrophages co-cultured with ovarian cancer stem-like cells through IL-8/STAT3 axis. *J Exp Clin Cancer Res*. 2019; 38(1):19. <https://doi.org/10.1186/s13046-018-1010-1>.

The original article can be found online at <https://doi.org/10.1186/s13046-018-1010-1>.

* Correspondence: Caojianguo2005@126.com; lx58616@163.com; tluox@126.com

[†]Yingxia Ning and Weifeng Feng contributed equally to this work.

⁴Department of Pharmaceutical Science, Medical College, Hunan Normal University, Changsha 410013, China

³Department of preclinical medicine, Medical College, Hunan Normal University, Changsha 410013, China

²The First Affiliated Hospital of Jinan University, Guangzhou 510632, China

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 9 Combination of GEN and STAT3 shRNA inhibited xenograft growth by co-injection of SKOV3-derived OCSLCs and THP-1 macrophages. The nude mouse xenograft model using co-injection with OCSLCs/THP-1 macrophages was treated with GEN (50 mg/kg) and Ad-shSTAT3 alone or in combination. The size **(a)** volume **(b)**, weight **(c)**, histological examination (HE staining) and the expression of CD68, IL-8 and p-STAT3 (immunohistochemical staining) **(d)** of xenografts were shown (scale bar, 100 μ m). * $P < 0.05$, vs the model control group; ** $P < 0.05$, vs treated with GEN (50 mg/kg) or Ad-shSTAT3 alone (means \pm SD, $n = 4$)