CORRECTION Open Access

Correction to: Pre-metastatic niche triggers SDF-1/CXCR4 axis and promotes organ colonisation by hepatocellular circulating tumour cells via downregulation of Prrx1



Yujun Tang^{1†}, Yishi Lu^{2,3†}, Yuan Chen^{1†}, Lei Luo^{1†}, Lei Cai¹, Bangjian Peng⁴, Wenbin Huang¹, Hangyu Liao¹, Liang Zhao^{2,3,4*} and Mingxin Pan^{1*}

Correction to: J Exp Clin Cancer Res 38, 473 (2019) https://doi.org/10.1186/s13046-019-1475-6

Following publication of the original article [1], the authors identified minor errors in image-typesetting in Fig. 3, specifically:

- Figure 3A: shNC image replaced with correct image
- Figure 3D: Nanog western blot replaced with correct image

The corrected figure is provided here. The corrections do not have any effect on the results or conclusions of the paper. The original article has been corrected.

Author details

¹Second Department of Hepatobiliary Surgery, Zhujiang Hospital, Southern Medical University, Guangzhou, China. ²Department of Pathology, Nanfang Hospital, Southern Medical University, Guangzhou, China. ³Department of Pathology, School of Basic Medical Sciences, Southern Medical University, Guangzhou, China. ⁴Department of Hepatobiliary Surgery, the Fifth Affiliated Hospital, Southern Medical University, Guangzhou, China.

Published online: 16 July 2021

Reference

 Tang Y, Lu Y, Chen Y, Luo L, Cai L, Peng B, et al. Pre-metastatic niche triggers SDF-1/CXCR4 axis and promotes organ colonisation by hepatocellular circulating tumour cells via downregulation of Prrx1. J Exp Clin Cancer Res. 2019;38(1):473. https://doi.org/10.1186/s13046-019-1475-6.

The original article can be found online at https://doi.org/10.1186/s13046-019-1475-6.

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*} Correspondence: liangsmu@foxmail.com; pmxwxy@sohu.com

[†]Yujun Tang, Yishi Lu, Yuan Chen and Lei Luo contributed equally to this

²Department of Pathology, Nanfang Hospital, Southern Medical University, Guangzhou, China

¹Second Department of Hepatobiliary Surgery, Zhujiang Hospital, Southern Medical University, Guangzhou, China




Fig. 3 Loss of Prrx1 is essential for SDF-1-induced stemness and migratory potential of HCC cells. **a** Migrating cells were counted under a microscope in five randomly selected fields. Bars represent number of migrated cells. **b** Drug resistance (epirubicin) was evaluated in indicated cells by CCK-8 assay. **c** Phase-contrast images of sphere-forming assays of indicated cells. **d** Western blotting and RT-qPCR were used to detect protein and mRNA expression of stemness markers, respectively. **e** Invading cells were counted under a microscope in five randomly selected fields. Bars represent number of migrated cells. **f** Drug resistance (epirubicin) was evaluated in indicated cells by CCK-8 assay. **g** Phase-contrast images of sphere-forming assays of indicated cells. **h** Western blotting was performed to detect the expression of stemness markers. * *P* < 0.05, *** *P* < 0.01