CORRECTION Open Access



Correction: Mechanical stress shapes the cancer cell response to neddylation inhibition

Frédérique Mittler¹, Patricia Obeïd¹, Vincent Haguet¹, Cédric Allier², Sophie Gerbaud¹, Anastasia V. Rulina^{1,3}, Xavier Gidrol¹ and Maxim Y. Balakirev^{1*}

Correction to: J Exp Clin Cancer Res 41, 115 (2022) https://doi.org/10.1186/s13046-022-02328-y

Following publication of the original article [1], an error was identified in Fig. 1; specifically:

• Fig. 1b: Incorrect image used for Pulse-MLN cells (bottom right panel); the correct image is now used

The corrected figure is given here. The correction does not have any effect on the final conclusions of the paper. The original article has been corrected.

Author details

¹University Grenoble Alpes, CEA, INSERM, IRIG, Biomics, 38054 Grenoble, France. ²University Grenoble Alpes, CEA, LETI, 38054 Grenoble, France. ³University of Bergen, Bergen, Norway.

Published online: 06 May 2022

Reference

 Mittler F, Obeïd P, Haguet V, et al. Mechanical stress shapes the cancer cell response to neddylation inhibition. J Exp Clin Cancer Res. 2022;41:115. https://doi.org/10.1186/s13046-022-02328-y.

The original article can be found online at https://doi.org/10.1186/s13046-022-02328-y.

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visithttp://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*}Correspondence: maxim.balakirev@cea.fr

¹ University Grenoble Alpes, CEA, INSERM, IRIG, Biomics, 38054 Grenoble, France

Mittler et al. J Exp Clin Cancer Res (20.




Fig. 1 Neddylation inhibition induces distinct phenotypes in PCa cells. **A, B** Effect of 100 nM MLN on PCa colony growth in soft agar ("steady"). In the "pulse" regimen, the cells were pretreated with 100 nM MLN for 1 h before seeding (see also Supplementary Figure S1A). The histograms show the colony size distribution of control (black)- and MLN (red)-treated cells. The numbers indicate the total area occupied by the colonies of the given size. Scale bar = 1 mm. **C, D** Wound healing assay with LNCaP (**C**) and PC3 (**D**) cells (mean \pm) S.D., n = 5 for LNCaP, n = 6 for PC3, ***-p < 0.001). 100 nM MLN was added just before monolayer scratching. Scale bar = 500 μ m (**E, F)** Effect of 100 nM MLN on LNCaP (**E**) and PC3 (**F**) tumoroid growth over 10 days. Scale bar = 150 μ m. In all experiments with MLNs, DMSO was used as a vehicle control